INTEGRATED MANAGEMENT OF CUCUMBER MOSAIC VIRUS OF CHILLI (CAPSICUM ANNUM L.)

M. S. Rahman^{1*}, K. Jahan¹, R. Islam¹, K. M. Alam¹, M. R. Karim² and M. M. Rahman¹

¹Plant Pathology Division, Bangladesh Agricultural Research Institute, Gazipur ²Pomology Division, Horticulture Research Centre, BARI, Joydebpur, Gazipur ^{*}Corresponding author: mdsiddiqurrahman@yahoo.com

ABSTRACT

Rahman, M. S., Jahan, K., Islam, R., Alam, K. M., Karim, M. R. and Rahman, M. M. 2022. Integrated management of cucumber mosaic virus of chilli (*Capsicum annum*). Bangladesh J. Plant Pathol. 38(1&2): 21-26.

Cucumber mosaic virus (CMV) is one of the destructive virus disease causing serious yield and quality loss in chilli worldwide. The experiment was conducted at the research field of Plant Pathology Division, Bangladesh Agricultural Research Institute during two consecutive years to find out the effective management option against CMV of chilli. Six packages with different combinations of seed bed netting, using yellow trap, spraying of Imidacloprid, bio-neem, and paraffin oil were tested against CMV of chilli in field condition. Significant variation of disease incidence and severity was found in different treatment packages. Disease incidence ranged from 10.95 to 38.50 % and the severity varied from 0.93 to 3.50 in different treatment packages. The lowest disease incidence (10.95%) and severity (0.93) was found in treatment package I (Net in seed bed + one spray of Imidacloprid 0.1% at 5 days before

transplanting + Sticky yellow trap in the plot + 3 sprays of Imidacloprid 0.1% at 15 days interval starting after observing the vector population by the yellow trap). The package II (Same as T1 except use Bio-Neem 0.2 % instead of Imidacloprid) also showed lower disease incidence (12%) and severity (1.10). The highest yield (12.24 t/ha) was recorded from package I followed by package II (12.00 t/ha). The reduction of disease incidence was 71.56 % in package I and 68.83 % in package II and thereby increased crop yield by 64.52% and 61.29 %. The package I and package II were considered as the effective management option on the basis of minimum disease incidence, higher yield and marginal benefit cost ratio (1: 4.64 & 1:4.06). These management packages could be used for the management of CMV disease of chilli.

Key words: Integrated management, cucumber mosaic virus, chilli, capsicum annum

INTRODUCTION

Chilli (*Capsicum annuum* L.), belongs to the family Solanaceae is an important vegetable and spices crop having immense commercial importance. It is one of the most important crops grown worldwide and cultivated in the Indian sub-continent for vegetable, spices and industrial purposes (Kumar 2012). Chilli, particularly hot peppers, is an important crop in tropical and subtropical Asia, not only because of popular consumption but also their high market value and export potential. Due to worldwide distribution and polyphagous vectors the *Cucumber mosaic virus* (CMV) is one of the most important virus infecting vegetables worldwide (Palukaitis and Garcia-Arenal 2003). Disease, particularly those caused by viruses are considered the major constraints to economic production of peppers and yield losses ranging from 60-100 % in case of early infection (Singh and Cheema 1989, Shukor *et al.* 1989). The CMV is most abundant and appeared as a major devastating virus in the leading chilli growing regions of Himachal Pradesh in India (Biswas *et al.* 2013)).

In Bangladesh CMV is one of the most important viruses which cause serious yield losses in chilli and can reduce fruit yield up to 80% (Rahman 2008). It is spread naturally by aphid in a nonpersistent manner. Plant viruses are usually very difficult to control. Spraying of chemical insecticides

²⁰²² Bangladesh Phytopathological Society

is the only option available to the farmers for the management of CMV through vectors control. As the CMV is a non-persistence virus, only insecticidal spray may not control the disease effectively because of very short time is required to transmit the virus and vector is able to transmit virus before their death. Moreover, dependence on a single method is highly vulnerable and sometimes become failure to the management of CMV in chilli (Lepidot et al. 2001). Hence, the present study was undertaken to find out a suitable management option against Cucumber mosaic virus (CMV) of chilli.

MATERIALS AND METHODS

The experiment was conducted in the research field of Plant Pathology Division, Bangladesh Agricultural Research Institute (BARI), Joydebpur, Gazipur during two consecutive years using randomized completely block design (RBD) with three replications. The unit plot size was $2.5 \times 2.5 \text{ m}^2$. Six management packages with a control were used as treatment. The management packages were package I : Net in seed bed + one spray of Imidacloprid 0.1% at 5 days before transplanting + Sticky yellow trap in the plot + 3 sprays of Imidacloprid 0.1% at 15 days interval starting after observing the vector population (aphids) by the yellow trap, package II: Same as package I except spray with 0.2 % Bio-neem instead of Imidacloprid, package III: Same as package I except spray with 2% mineral oil (Paraffin oil), package IV: Net in seed bed + one spray of Imidacloprid 0.1% at 5 days before transplanting + 3sprays of Imidacloprid 0.1% at 15 days interval starting at 20 days after transplanting, package V: Same as package IV except spray with 0.2 % Bioneem instead of Imidacloprid, package VI: Same as package IV except spray with 2% mineral oil and package VII: Untreated control. The chilli variety BARI-Marich-1 was used in this experiment. Seedlings were raised in the nursery bed. Five weeks old seedlings were transplanted in the experimental field with a spacing of 50 \times 50 cm. Irrigation, drainage, weeding and other intercultural operations were done as and when required. The experimental plots were monitored regularly to observe CMV symptoms. The virus CMV was confirmed by the DAS-ELISA. The diseases incidence was calculated by using the following formula.

Disease incidence = $\frac{\text{No. of infected plant}}{\text{Total plants in the plot}} \times 100$ The severity of Cucumber mosaic virus (CMV) was determined according to Monma and Sakata (1997) based on a 0-4 scale as follows: Severity Index= $\frac{\Sigma(\text{Symptom index}\times\text{No. plant of sumptom index})}{\text{Total number of plants}}$

where the symptom index was as 0 = No Symptom, 1= Mild Mosaic, 2 = Mosaic, 3 = Mosaic & deformed leaf, and 4 = Mosaic and stunted plants.

Data on fresh fruit yield (kg/plot) were recorded and finally converted into fruit yield of chilli into ton ha-1. The data were analyzed statistically for analysis of variance (ANOVA) using MSTAT-C program and means were compared according to Duncan's Multiple Range Test (Gomez and Gomez 1984). The percent data were transformed using Arcsine transformation method before statistical analysis.

Economic analysis

Economic analysis was performed by partial budget technique as described by Rahman et al. (2011) following the formulae of economic analysis. Variable Cost = Cost (Taka) that vary in different packages; Gross Return (TR) = Yield in terms of money (@ Taka 20/kg (Yield price); Gross margin = Gross Return-Variable cost; Marginal benefit = Gross margin (packages) - Gross margin (control). The marginal benefit cost ratio (MBCR) was

calculated by the following formula: MBCR (over control) = $\frac{\text{Marginal benefit}}{\text{MBCR}}$

RESULTS AND DISCUSSION

Incidence and severity of Cucumber mosaic virus (CMV) disease of chilli

The incidence and severity of Cucumber mosaic virus disease in different management packages varied significantly (Fig. 1). The percent disease incidence ranged from 10.95 to 38.50 % and severity varied from 0.93 to 3.50 in different treatment packages. Incidence of CMV of all the management packages was lower as compared to untreated control. The highest disease incidence (38.50%) was found in control (package VII) and the lowest incidence (10.95 %) was recorded from package I which was statistically similar to package II (12%). The incidence of CMV in other packages ranged from 21 - 29% where package III (21%) and package IV (22.50%)were statistically similar. Similarly, the highest disease severity was found in package VII (control) and significantly lower disease severity of CMV were found in all the management packages over control. The lowest severity was found in package I, which was statistically similar to package II. However, treatments involving netting seedling, 1 spray of Imidacloprid 0.1% or Bio-neem 0.2 % at 5 days before transplanting and 3 sprays of Imidacloprid or Bio-neem at 15 days interval and use sticky yellow trap (package I and package II) were found very much effective as compared to other management packages in respect of disease incidence and severity (Fig. 1). It might be due to better control of CMV vectors (aphids) in the treated plots by netting seedlings and use of sticky yellow trap in the field, the seedlings remain disease free at early stage and spray of insecticides gave better control against CMV. The CMV, an aphid born non-persistent virus, could not be controlled by only insecticides pray as CMV vector required only few seconds to transmit virus from infected to healthy plant. So, the management packages i.e. use of disease-free seedling, sticky yellow trap, and spray insecticide effectively controlled the vectors and reduced the incidence as well as severity of CMV disease in chilli.

Fresh fruit yield of chilli

The fruit yield of chilli (BARI Marich-1) in different management packages varied significantly (Table 1). All the management packages gave higher fruit yield of chilli over control. The highest average fresh fruit yield of 7.65 kg/plot or 12.24 t/ha was found in the management package I (Net in seed bed + one sprav of Imidacloprid 0.1% at 5 days before transplanting + Sticky yellow trap + 3 sprays of Imidacloprid 0.1% at 15 days interval starting after observing the vector aphids by the yellow trap) which was statistically similar the package II (Netting seed bed + 1 spray of Bio-neem 0.2% at 5 days before transplanting + sticky yellow trap + 3 sprays of Bio-neem at 15 days' interval starting after observing the vector aphids by the yellow trap). The lowest fruit yield of 4.65 kg/plot or 7.44 t/ha was found in package VII (Untreated control). The fresh fruit yield of other management packages ranged from 5.00 kg/plot to 6.19 kg/plot or 8.00 t/ha to 9.90 t/ha.



Figure 1. Effect of different management packages on the incidence and severity of CMV disease of chilli

[Package I: Net in seed bed + one spray of Imidacloprid 0.1% at 5 days before transplanting + Sticky yellow trap + 3 sprays of Imidacloprid 0.1% at 15 days interval starting after observing the vector population (aphids) by the yellow trap, Package II: Same as package I except spray with 0.2 % Bio-Neem instead of Imidacloprid, Package III: Same as package I except spray with 2% mineral oil (Paraffin oil), Package IV: Net in seed bed + one spray of Imidacloprid 0.1% at 5 days before transplanting + 3 sprays of Imidacloprid 0.1% at 15 days interval starting at 20 days after transplanting, Package V: Same as package IV except spray with 0.2 % Bio-Neem instead of Imidacloprid 0.1% at 5 days before transplanting + 3 sprays of Imidacloprid 0.1% at 15 days interval starting at 20 days after transplanting, Package V: Same as package IV except spray with 0.2 % Bio-Neem instead of Imidacloprid, Package VI: Same as package IV except spray with 2% mineral oil, and Package VII: Untreated control].

All the management packages reduced the disease incidence of CMB in chilli and thereby increased fruit yield over control (Table 2). The average disease reduction ranged from 24.68% to 71.56% in different management packages. The highest reduction of disease incidence was 71.56% in management package I which was followed by management package II where disease reduction was 68.83% over control. The average increase of fresh fruit yield of chilli ranged from 7.53% to 64.52% in different management packages. The highest increase

of fresh fruit yield over control was recorded as 64.52% in package I which was followed by package II (61.29%). Other management packages reduced disease incidence from 24.68 to 45.45% and increased fresh fruit yield from 7.53 to 33.06%. The average fresh fruit yield of chilli was increased in different management packages might be due to decrease of disease incidence through vector control. These findings were in conformity with the previous findings of Rahman *et al.* (2020).

Management	Fresh fruit yield of chilli in two consecutive years					
packages	Fresh fruit yield (kg/plot)		Fresh fruit yield (t/ha)			
	1 st year	2nd year	Pooled	1 st year	2nd year	Pooled
Package I	7.70	7.60	7.65 a	12.32	12.16	12.24 a
Package II	7.20	7.80	7.50 a	11.52	12.48	12.00 a
Package III	6.05	6.32	6.19 b	9.68	10.12	9.90 b
Package IV	5.30	5.20	5.25 c	8.48	8.32	8.40 c
Package V	5.10	5.25	5.18 c	8.18	8.40	8.29 c
Package VI	4.85	5.15	5.00 c	7.76	8.24	8.00 c
Package VII	4.70	4.60	4.65 d	7.52	7.36	7.44 d
(Untreated Control)						
CV %			14.75			13.5

Table 1. Fruit yield of chilli in different management packages in two consecutive years

Means followed by same letters in row and column are not significantly different at 5% level by DMRT

Treatment packages	Disease Incidence		Reduction of	Yield	Yield increase	
	1 st year	2nd year	Pooled	disease incidence	(t/ha)	over control (%)
				over control (%)		
Package I	11.00	10.90	10.95 e	71.56	12.24 a	64.52
Package II	11.50	12.50	12.00 e	68.83	12.00 a	61.29
Package III	23.00	19.00	21 d	45.45	9.90 b	33.06
Package IV	25.00	20.00	22.50 d	41.56	8.40 c	12.90
Package V	23.00	27.00	25 c	35.06	8.29 c	11.42
Package VI	27.00	31.00	29.00 b	24.68	8.00 c	07.53
Package VII	37.00	40.00	38.50 a	-	7.44 d	-
(Untreated Control)						
CV %			14.75		13.5	

Table 2. Effect of different treatment packages on disease reduction and fruit yield of chilli

Means followed by same letters in row and column are not significantly different at 5% level by DMRT.

[Package I: Net in seed bed + one spray of Imidacloprid 0.1% at 5 days before transplanting + Sticky yellow trap + 3 sprays of Imidacloprid 0.1% at 15 days interval starting after observing the vector population (aphids) by the yellow trap, Package II: Same as package I except spray with 0.2 % Bio-Neem instead of Imidacloprid, Package III: Same as package I except spray with 2% mineral oil (Paraffin oil), Package IV: Net in seed bed + one spray of Imidacloprid 0.1% at 5 days before transplanting + 3 sprays of Imidacloprid 0.1% at 15 days interval starting at 20 days after transplanting, Package V: Same as package IV except spray with 0.2 % Bio-Neem instead of Imidacloprid 0.1% at 0.2 % Bio-Neem instead of Imidacloprid 0.1% at 15 days interval starting at 20 days after transplanting, Package V: Same as package IV except spray with 0.2 % Bio-Neem instead of Imidacloprid, Package VI: Same as package IV except spray with 2% mineral oil, and Package VII: Untreated control].

Economic analysis

The management packages of Cucumber Mosaic Virus (CMV) disease of chilli appreciably increased the gross return over control (Table 3 & 4). The gross return was highest in management package I (244800 Tk. ha⁻¹) followed by management package II (240000 Tk. ha⁻¹). The lowest gross return was obtained from the untreated Control (148800 Tk. ha⁻¹). The gross returns of other treatments ranged from 160000 to 198000 Tk. ha⁻¹ (Table 3). Marginal analysis pointed out that the management packages I, II & III considerably increase marginal benefit cost ratio (MBCR 4.64, 4.06 & 2.28) over control (Table

4). The highest MBCR was obtained with package I and the lowest with package VI. The results showed that additional investment of Taka 1 in package I over control had additional income of Taka 4.64 followed by package II, III & IV with Tk. 4.06, 2.28, and 1.56, respectively. The cost and return and MBCR from the economic analysis indicated that the management packages I, II, III, & IV were economically viable and maximum gain could be obtained from package I (Net in seed bed + one spray of Imidacloprid 0.1% at 5 days before transplanting + Sticky yellow trap + 3 sprays of Imidacloprid 0.1% at 15 days interval starting after observing the vector aphids by the yellow trap).

Table 3. Cost and return in different management packages of CMV disease of chilli

Packages	*Var. Cost (Tk ha ⁻¹)	Yield (t ha ⁻¹)	**Gross return (Tk ha ⁻¹)
Package I	17000	12.24	244800
Package II	18000	12.00	240000
Package III	15000	9.90	198000
Package IV	7500	8.40	168000
Package V	8000	8.29	165800
Package VI	7000	8.00	160000
Package VII (Untreated Control)	-	7.44	148800

* Var. Cost: Cost that vary in different packages; ** Whole Sell rate of green chilli @ TK 20.00/Kg

Packages	Gross return	Variable Cost	Gross margin	Marginal benefit	MBCR		
	(Tk ha ⁻¹)						
Package I	244800	17000	227800	79000	4.64		
Package II	240000	18000	222000	73200	4.06		
Package III	198000	15000	183000	34200	2.28		
Package IV	168000	7500	160500	11700	1.56		
Package V	165800	8000	157800	9000	1.13		
Package VI	160000	7000	153000	4200	0.60		
Package VII (Untreated Control)	148800	-	148800	-	-		

Table 4. Marginal analysis of different management packages of CMV disease of chilli

(MBCR: Marginal benefit cost ratio)

[Package I: Net in seed bed + one spray of Imidacloprid 0.1% at 5 days before transplanting + Sticky yellow trap + 3 sprays of Imidacloprid 0.1% at 15 days interval starting after observing the vector population (aphids) by the yellow trap, Package II: Same as package I except spray with 0.2 % Bio-Neem instead of Imidacloprid, Package III: Same as package I except spray with 2% mineral oil (Paraffin oil), Package IV: Net in seed bed + one spray of Imidacloprid 0.1% at 5 days before transplanting + 3 sprays of Imidacloprid 0.1% at 15 days interval starting at 20 days after transplanting, Package V: Same as package IV except spray with 0.2 % Bio-Neem instead of Imidacloprid 0.1% at 0.2 % Bio-Neem instead of Imidacloprid 0.1% at 15 days interval starting at 20 days after transplanting, Package V: Same as package IV except spray with 0.2 % Bio-Neem instead of Imidacloprid, Package VI: Same as package IV except spray with 2% mineral oil, and Package VII: Untreated control].

The findings of the present investigation revealed that incidence of Cucumber mosaic virus (CMV) disease of chilli was considerably reduced by the management package, i.e. net in seed bed + one spray of Imidacloprid 0.1% or Bio-neem 0.2 % at 5 days before transplanting + Sticky yellow trap + 3 sprays of Imidacloprid 0.1% or Bio-neem 0.2 % at 15 days interval starting after observing the vector aphids by the yellow trap. The management packages I and II were considered as effective management options on the basis of reduced disease incidence, higher fruit yield and marginal benefit cost ratio. Marginal cost benefit analysis indicated that the two management packages (I & II) were economically viable and cost effective. However, these management packages could be used as effective options for management of CMV disease of chilli in Bangladesh.

ACKNOWLEDGEMENTS

The author thankfully acknowledges the scientists and stuffs of plant pathology division, BARI for their assistance in data collection and also to SRC for supplying the seeds of BARI Marich-1.

LITERATURE CITED

- Biswas, K., Hallan, V., Zaidi, A.A. and Pandey, P.K. 2013. Molecular evidence of *Cucumber mosaic virus* subgroup II infecting *Capsicum annuum* L. in the Western region of India. Inter. J.of Current discoveries and innovations. 2 (2): 97-105.
- Gomez, K.A. and Gomez. A.A. 1984. Statististical Procedures for Agricultural Research. Second edn. John. Wiley and Sons. Inc. New York. pp. 304-307.
- Kumar, R.V., Singh, A.K. and Chakraborty, S. 2012. A new monopartite begomovirus species, Chilli leaf curl Vellanad virus, and associated beta-satellites infecting chilli in the Vellanad region of Kerala, India. New Dis. Rep. 25: 20.
- Lepidot, M. Friedmann, M., Pilowsky, M., Ben-Joseph, R. and Cohen, S. 2001. Effect of host plant resistance to tomato yellow leaf curl virus (TYLCV) on virus acquisition and transmission by its whitefly vector. Phytopathology. 91: 1209-1213.

- Monma, S., and Sakata, Y. 1997. Screening of *Capsicum* accessions for resistance to *Cucumber mosaic virus*. J. Japan Soc. Hortic. Sci.65:769-776.
- Palukaitis, P. and Garcia-Arenal, F. 2003. "Cucumoviruses," Advances in Virus Research. 62: 241-323.
- Rahman, M. S. Ahmed, A. U. Jahan, K. Khatun, F. 2020. Management of Cucumber mosaic virus (CMV) infecting cucumber in Bangladesh. Bangladesh J. Agril. Res. 45 (1): 65-76. DOI: https://doi.org/10.3329/bjar.v44i4.45693.
- Rahman, M.H., Islam M.R., Jahiruddin, M. and Haque, M.Q. 2011. Economics of fertilizer use in the Maize-Mungbean/Dhaincha-T.aman rice cropping pattern. J. Bangladesh Agril. Univ. 9(1): 37–42.
- Rahman, M.S. 2008. Performance of chilli varieties against anthracnose and major viral diseases.A MS thesis submitted to the Plant Pathology Department, BSMRAU, Gazipur. P 56-59.
- Shukor, N.M., Chew, B.H. Noraini, S. and Roff, M.N.M. 1989. Tomato and chilli pepper production in Malaysia. P. 508-520. In. S.K. Green, T.D. Griggs and B.T. McLean eds. Tomato and Pepper Production in the Tropics. Proceedings of the International Symposium on Integrated Management Practices. Tainan, 21-26 march, Asian Vegetable Research and Development Center, Taiwan.
- Singh, J.H. and Cheema, D.S. 1989. Present status of tomato and pepper production in the tropics. P. 452-471. In. S.K. Green, T.D. Griggs and B.T. McLean eds. Tomato and Pepper Production in the Tropics. Proceedings of the International Symposium on Integrated Management Practices. Tainan, 21-26 march, Asian Vegetable Research and Development Center, Taiwan.